

Nonparametric Stability Analysis of Starch Content of Gamma Irradiated Cassava at Three Locations in West Java, Indonesia

Ferra Anggita Agustina^A, Nurul Khumaida^B, Sintho Wahyuning Ardie^B, and Muhamad Syukur^{*B}

^A Department of Agronomy and Horticulture, Faculty of Agriculture, IPB University, Bogor 16680, Indonesia

^B Department of Agronomy and Horticulture, Faculty of Agriculture, IPB University, Bogor 16680, Indonesia

*Corresponding author; email: muhsyukur@apps.ipb.ac.id

Abstract

Cassava is one of the largest starch producing tuber crops in Indonesia. Tapioca from cassava starch can be used as a raw materials for a variety of foods including biscuits, instant porridge, meatball, sausage, nuggets, and flour condiment. Breeding programs to improve cassava yield and starch content by using mutagen gamma rays irradiation was tested in four cassava generations (M_1V_4). Cassava is propagated by stem cuttings and the new characteristic obtained from gamma irradiation mutation is stable and can be passed from one generation to the next. Cassava mutants were obtained by evaluating the performance in different environmental conditions. The testing of yield adaptability and stability through a series of multi location is an important step before a new variety can be released. The aim of this research was to compare nonparametric stability and to evaluate the stability of the starch content of 16 genotypes (14 mutants and 2 varieties) in three locations in West Java Province i.e. Tapos (Depok), Cikarawang (Bogor), and Ciseeng (Bogor). Experiments were conducted in a completely randomized block design with three replications nested in each environment. Testing of yield stability in this study used four approaches with 10 parameters of nonparametric method. Starch content was estimated using gravimetric method. Based on frequency stability ranking of starch content from gravimetric method the G63142 genotype had the highest starch content estimation (29.99%), and the top five genotypes with high starch content were G63142, G61142, "Manggu", G62151, and G63124. G61142 was categorized as genotype with static and dynamic stability, therefore this genotype is a very potential mutant to be released, whereas G63124 is categorized as a genotype with a static stability. Genotypes G63142, G62151, and "Manggu" had variable but relatively high starch content.

Keywords: gravimetric, mutant, static stability, dynamic stability

Introduction

Cassava is one of the important carbohydrate producing crops after wheat, rice, corn, potato, and barley (Lebot, 2009). FAO (2017) showed that Indonesia ranks as the fourth largest cassava producing country after Nigeria, Thailand, and Brazil in 2016. Cassava is used as raw material for food and non-food industries, including feed, flour, and fuel.

Cassava contributed the largest percentage of starch from tuberous crops (29%), followed by sweet potato (26%) and potato (5%). Cassava starch has granule form which are round with a truncated end; so it is different from corn starch which is hexagonal, and sago starch which is oval (Murtiningrum, 2012). Tapioca from cassava starch can be made into biscuits, instant porridge, processed foods such as meatball, sausage, nuggets, and flour condiment. Hydrolysis of modified cassava flour can be processed into soft drink, milk, and ketchup (Haloho, 2014).

Indonesian cassava starch export value in 2017 was US\$ 5.60 billion, whereas the imports value was US\$ 126.4 billion (BPS, 2017). The Indonesian cassava starch exports decreased to US\$ 5.28 billion in 2018, whereas the import value increased to US\$ 185.6 billion at the same period. Even though Indonesia has exported cassava starch and chips, the national production faces uncertainties of productivity because of pest outbreak and weather fluctuations.

Efforts to obtain new cassava varieties through conventional breeding have limitations when using stem cuttings as the genetic material. Cassava is usually propagated vegetatively because it only flowers and produce seeds at 800 m above sea level, therefore cassava has a relatively low genetic variance. Vegetative propagation through stem cuttings has many advantages; the new characteristic obtained from gamma irradiation are stable and can be passed

from one generation to the next. Genetic variance can be increased by gene recombination, hybridization, genetic manipulation, mutation induction, or polyploidy (Syukur et al., 2012). Breeding through mutation induction on cassava has been used to develop new superior varieties; Fahreza (2014) and Maharani (2015) have reported an improved yield and starch content in cassava through mutation breeding using mutagen gamma ray irradiation.

According to the regulation by Indonesian Ministry of Agriculture (2011) the new crop genotypes have to be characterized for yields, adaptability to different environments and locations prior. Different genotypes might express different characters in different environments (Harahap, 1982).

This study evaluated yield stability of different genotypes of cassava at three locations in West Java Province. The locations were selected based on the ideal topology and topography for growing cassava. Genotype performance might change in different environments known as interaction of genotype and environment ($G \times E$). Interaction of genotype and environment is very important when a mutant genotype is to be grown in different environments. Baye et al (2011) stated the importance of developing varieties that are specifically adapted to a particular environment (Baye et al., 2011).

Yield stability studies based on the interaction of genotype \times environment have been widely conducted, e.g. Finlay-Wilkinson (1963), Eberhart-Russell (1966), Francis and Kannenberg (1978), and Gauch (2006). The interaction of genotype and environment can also hinder the progress of selection, disrupt the selection of excellent varieties in a testing of varieties and difficult to make appropriate conclusion if a genotype test is performed in a wide range of environments. The testing of yield stability in this study used four approaches with 10 parameters of nonparametric method. Huehn (1990) suggests that the nonparametric procedure has several advantages compared to the parametric stability. These benefits are the reduction of bias caused by outliers, no assumptions needed of the observed values and ease of use in interpreting, adding, or deleting unsuitable genotypes.

The starch content of cassava tubers was usually measured using a gravimetric method. Subekti et al. (2018) evaluated starch content of cassava "Gajah", a mutant derived from mutation induction using gamma ray irradiation. The value of gravimetric method was included to the formula and the percent of starch content was measured. The starch content estimation (expressed in %) from gravimetric method

had similar results with laboratory test.

The aim of this research was to compare non parametric stability measures and to evaluate the starch content stability of 16 genotypes of cassava that consists of 14 mutants and two varieties, in three locations in West Java Province i.e. Depok (Tapos), Bogor (Cikarawang), and Bogor (Ciseeng).

Materials and Methods

The genetic materials used for this experiment were 14 putative mutant lines (G61122, G61142, G61143, G61151, G62143, G62132, G62151, G62153, G631111, G63121, G63124, G63142, G63143, G63153) and two national varieties ("Gajah" and "Manggu"). The putative mutants were selected from the previous generation (Subekti et al., 2018). Breeding programs to improve cassava yield and starch content was conducted using mutagen gamma ray irradiation with five doses (0, 15, 30, 45, 60 Gy), the crop performance was evaluated in four generations. Based on the regulation by Indonesian Ministry of Agriculture (2011), genotype performances have to be stable in different environment.

The research was conducted at Cikarawang (Bogor) with altitude of 192 m above sea level, Ciseeng (Bogor) with altitude of 125 m above sea level, and Tapos (Depok) with altitude of 110 m above sea level. The experiment was conducted in a randomized complete block design with three replication in each location. Plant maintenance involved regular watering, fertilizing, and weeding. Pest and disease control was conducted only when required. Gravimetric analysis on fresh tubers was conducted twice at harvest. Cassava tubers were harvested 11 months after planting.

Starch content was analyzed using a gravimetric method by Kawano et al. (1987). Tubers with length of > 20 cm and diameter of >4 cm were weighed in the air (x) and in the water (y). Specific gravity was calculated by the formula $(x/x-y)$. The value of specific gravity was included to the formula $(112.1 \times SG) - 106.4$ to get the percentage of starch content.

Nonparametric stability analysis was based on ranks classification (Ackura et al., 2006). Akcura et al. (2006) showed that nonparametric procedure has an advantage of reducing bias caused by outliers. Rahadi et al. (2013) used the nonparametric stability analysis based on two different concepts, static and dynamic stability, of yield for nine chili pepper genotypes in eight environments.

Data analysis consisted of the analysis of variance followed by nonparametric stability analysis with 10 methods of S_i^1 , S_i^2 , S_i^3 , and S_i^6 (Nassar and Huehn, 1987), RS (Kang, 1988), Top (Fox et al., 1990), and NPI^1 , NPI^2 , NPI^3 and NPI^4 (Thennarasu, 1995) against the production characteristics of each plant using the following formula (Syukur et al., 2012) :

$$S_i^{(1)} = 2 \sum_j \sum_{j'=j+1}^n |r_{ij} - r_{ij'}| / [n(n-1)]; S_i^{(2)} = \sum_{j=1}^n (r_{ij} - \bar{r}_i)^2 / (n-1); S_i^{(3)} = \frac{\sum_{j=1}^n (r_{ij} - \bar{r}_i)^2}{r_i}; S_i^{(6)} = \frac{\sum_{j=1}^n |r_{ij} - \bar{r}_i|}{r_i};$$

$$NP_i^{(1)} = \frac{\sum_{j=1}^n |r_{ij}^* - M_{di}^*|}{n}; NP_i^{(2)} = \frac{(\sum_{j=1}^n |r_{ij}^* - M_{di}^*| / M_{di}^*)}{n}; NP_i^{(3)} = \frac{\sqrt{\sum_{j=1}^n (r_{ij}^* - \bar{r}_i^*)^2 / n}}{r_i}; NP_i^{(4)} = \frac{2}{n(n-1)} \sum_j \sum_{j'=j+1}^n |r_{ij}^* - r_{ij'}^*| / \bar{r}_i$$

where n is the number of location, r_{ij} is genotype ranks-i in environments-j, r_{ij}^* is the rank of the genotype from a corrective values, M_{di}^* is median from corrective value, and M_{di} is median from all environments.

Result and Discussion

Interaction of Genotype and Environment (GxE)

Starch content from fresh tubers were analyzed by the combined variant analysis. Cassava genotypes have different percentage of starch content, whereas the environment and interaction of genotype x environment had no significant effect on starch content (Table 1), therefore each genotype could perform equally well at the three locations.

Subekti et al. (2018) reported that in terms of yield stability, the M1V4 generation mutants had stable yields, however different mutants were different in their starch content. Anasari et al. (2017) showed that there was no significant interaction of genotype x environment in affecting Pakchoy yields in three different locations, and the genotypes tested had high diversity with high adaptability to each location.

Figure 1 showed the interaction response indicated by the starch content fluctuations at each location. A

high diversity of macro environment is equal with high environment diversity for each genotypes (Satoto et al. (2009). Three cassava genotypes, G61142, G61143, and G62132, had similar starch content at every location.

Chahal and Gosal (2002) showed that the qualitative




Figure 1. Genotype x environment interaction (GxE) of 16 cassava genotypes at three geographical locations in West Java, Indonesia

interaction of genotypes were identified by yield frequency ranking in each location. Based on frequency ranking in three locations, some genotype were stable only on two locations (Table 2). G63142 had the highest starch content estimation (29.99%), whereas G62132 had the smallest starch content estimation (18.22%). Both showed good performance in Depok and Cikarawang. The last generation was tested by Subekti et al. (2018) and 23.86% had the highest and 12.60% had the smallest starch

Table 1. Variant analysis of the combined starch content (%) from three geographic locations

Sources of variants	DF	SS	MS	F calc	Pr > F
Environment (E)	2	157.4353	78.7176	3.84	0.1123 ^{ns}
Replication/Environment	3	43.9847	14.6615	1.31	0.5105 ^{ns}
Genotype (G)	15	832.5972	55.5064	2.26	0.0281*
G x E	30	737.5985	24.5866	1.31	0.2020 ^{ns}
Errors	45	844.1654	18.7592		
Total	95	2648.249			

Note: *= significantly different; ns= not significantly different at $\alpha=5\%$

Table 2. Starch content estimation (%) of 16 cassava genotypes at three geographical locations in West Java, Indonesia

No	Genotype	E1	Rank	E2	Rank	E3	Rank	Average
		-----%-----						
1	G61122	24.28	9	22.25	8	13.76	16	21.10cd
2	G61142	23.18	10	27.67	3	25.51	3	25.45abc
3	G61143	19.00	15	21.41	11	19.81	13	20.07cd
4	G61151	21.17	13	21.08	13	21.49	8	21.25bcd
5	G62143	21.68	11	21.47	10	25.76	1	22.97bcd
6	G62132	16.93	16	17.49	16	20.25	12	18.22d
7	G62151	29.40	4	26.67	4	21.23	9	25.77abc
8	G62153	20.54	14	20.53	15	18.83	15	19.97cd
9	G63111	21.20	12	23.14	6	19.76	14	21.36bcd
10	G63121	24.44	8	21.48	9	25.76	2	23.97bcd
11	G63124	32.66	2	23.27	5	24.61	6	25.72abc
12	G63142	39.29	1	29.75	2	20.93	10	29.99a
13	G63143	25.29	7	21.27	12	19.81	4	23.93bcd
14	G63153	30.05	3	20.70	14	20.26	11	23.67bcd
15	“Gajah”	28.12	5	22.28	7	25.20	5	25.20abc
16	“Manggu”	27.06	6	29.79	1	21.92	7	26.25ab
Environment average		25.27		23.14		21.89		23.77

Note: E1= Depok (Tapos), E2= Bogor (Cikarawang), E3= Bogor (Ciseeng); values followed by the same letter within the same column was not significantly different based on Duncan's test at $\alpha=5\%$.

content estimation. These genotypes demonstrated an increase in starch content in the three locations compared to those in the previous location.

Nonparametric Stability Analysis of Starch Content at Three Locations in West Java, Indonesia

The testing of yield adaptability and stability through a series of multi location evaluation is an important step before a new variety is released (Syukur et al., 2012). Stability studies based on Gx E interaction need to be complemented with nonparametric methods to reduce bias caused by outliers. In addition, this method had no assumption of the values, easy to interpret and to add, or to delete unsuitable genotypes without destroying the population (Huehn, 1990).

De Vita et al. (2010) suggested that the released genotypes should have high adaptability and high stability. Mohammadi et al. (2010) separated the different methods based on two concepts of stability, dynamically stable (adaptation followed the environmental index) and statically stable (yield stable in each environment). The farmers need stable and high yielding genotypes categorized by dynamic stability (Zulhayana, 2010). The stability values obtained from the Nassar and Huehn (1987) method

showed that G63142 genotype had the biggest value with 29.99% starch content estimation (Table 3), followed by G61142 and local genotype “Manggu”. The smallest starch content estimation was found in G62132 (18.23%). Sabaghnia et al. (2006) showed the smallest parametric stability genotype was more stable compared to the other genotypes. Based on $S1^{(1)}$ parameter of stability index (Nassar and Huehn, 1987) “Gajah” and G61142 genotype were stable, whereas $S1^{(2)}$, $S1^{(3)}$ and $S1^{(6)}$ parameter showed that G62153 genotype was stable. “Gajah” was the most stable compared to the local genotype “Manggu” in each environment. Fox et al. (1980) identified that genotypes in the top ranks from each environment were categorized as stable.

Table 3 shows that three cassava mutants, G61142, G62151, and G63142, had the highest ranks in TOP layer, therefore these mutants are recommended and adaptive. Rank-sum method by Kang (1988) was the sum of ranking in starch content estimation (%) obtained by Shukla method. The smallest value indicates a stable genotype, and G61142 was the most stable genotype. Table 4 shows all data from 10 parameters of 16 cassava genotypes in three environments.

Table 3. Average starch content estimation and nonparametric stability values of 16 cassava genotypes in three locations

Geno	SCE (%)	S1 ⁽¹⁾	S1 ⁽²⁾	S1 ⁽³⁾	S1 ⁽⁶⁾	TOP	MID	LOW	RS	NP _i ¹	NP _i ²	NP _i ³	NP _i ⁴
G61122	20.10	10.67	17.33	14.81	5.08	0.00	33.33	66.67	27	4.33	0.43	0.52	0.49
G61142	26.95	3.00	3.00	5.25	3.25	66.67	33.33	0.00	3	1.33	0.44	0.47	0.44
G61143	20.08	5.67	1.33	1.04	1.45	0.00	0.00	100.00	21	3.33	0.30	0.36	0.32
G61151	21.22	6.00	8.33	6.54	3.63	0.00	33.33	66.67	15	2.33	0.18	0.25	0.22
G62143	22.97	9.67	50.33	48.67	8.55	0.00	0.00	100.00	27	5.00	0.50	0.72	0.64
G62132	18.23	6.67	5.33	4.04	2.85	33.33	0.00	66.67	23	4.33	0.27	0.37	0.32
G62151	25.77	7.00	10.33	9.74	4.35	66.67	33.33	0.00	11	3.33	0.83	0.84	0.79
G62153	19.97	5.67	0.33	0.56	1.02	0.00	0.00	100.00	16	1.33	0.09	0.11	0.09
G63111	21.37	10.00	17.33	24.60	6.31	0.00	33.33	66.67	16	3.33	0.28	0.41	0.38
G63121	24.04	5.33	16.33	14.16	5.37	33.33	0.00	66.67	16	3.33	0.37	0.61	0.50
G63124	25.37	3.33	1.00	1.20	1.20	33.33	66.67	0.00	8	2.00	0.40	0.50	0.44
G63142	29.99	9.33	24.33	23.97	6.97	66.67	0.00	33.33	17	5.00	2.50	1.41	1.18
G63143	23.93	6.67	16.00	18.00	4.50	33.33	33.33	33.33	16	3.00	0.38	0.47	0.42
G63153	23.67	11.67	39.00	74.44	12.22	33.33	0.00	66.67	21	4.67	0.42	0.65	0.59
“Gajah”	25.20	3.00	1.33	2.30	2.12	0.00	100.00	0.00	21	3.33	0.67	0.79	0.75
“Manggu”	26.26	6.33	12.00	20.80	6.40	33.33	66.67	0.00	14	4.00	0.57	1.00	0.89

Note: SCE= starch content estimation per plant (%); S1⁽¹⁾, S1⁽²⁾, S1⁽³⁾, S1⁽⁶⁾ (Nassar and Huenh, 1987); Top (Fox et al., 1990); RS= Rank-sum by Kang (1988); NP_i¹, NP_i², NP_i³, and NP_i⁴ by Thennarasu (1995).

Table 4. Ranking of starch content estimation (%) and nonparametric stability of 16 cassava genotypes in three locations

Geno	SCE (%)	S1 ⁽¹⁾	S1 ⁽²⁾	S1 ⁽³⁾	S1 ⁽⁶⁾	TOP	RS	NP _i ¹	NP _i ²	NP _i ³	NP _i ⁴
G61122	13	15	12	10	10	12	16	12	10	9	9
G61142	2	1	5	6	6	2	1	1	11	7	8
G61143	14	5	3	2	3	12	10	6	12	3	4
G61151	12	7	7	7	7	12	5	4	2	2	2
G62143	10	13	16	15	15	12	16	15	5	12	12
G62132	16	9	6	5	5	7	12	13	3	4	3
G62151	4	11	8	8	8	2	3	7	15	14	14
G62153	15	6	1	1	1	12	6	2	1	1	1
G63111	11	14	13	14	12	12	6	8	4	5	5
G63121	7	4	11	9	11	7	6	9	6	10	10
G63124	5	3	2	3	2	7	2	3	8	8	7
G63142	1	12	14	13	14	2	7	16	16	16	16
G63143	8	10	10	11	9	7	6	5	7	6	6
G63153	9	16	15	16	16	7	10	14	9	11	11
“Gajah”	6	2	4	4	4	12	10	10	14	13	13
“Manggu”	3	8	9	12	13	7	4	11	13	15	16

Note: SCE= starch content estimation per plant (%); S1⁽¹⁾, S1⁽²⁾, S1⁽³⁾, S1⁽⁶⁾ (Nassar and Huenh 1987); Top (Fox et al. 1990); RS= Rank-sum by Kang (1988); NP_i¹, NP_i², NP_i³, and NP_i⁴ by Thennarasu (1995).

Syukur et al. (2012) showed that $S1^{(1)}$ and $S1^{(2)}$ parameter were placed in the same rank, whereas these parameters showed different values (Table 3). Only two genotypes, G61151 and G63143, were placed in the same rank. This shows that the tested genotypes behave similarly in different environment, whereas $S1^{(3)}$ and $S1^{(6)}$ parameters combined the yield and stability in each environment.

Nine genotypes (G61122, G61142, G61151, G62143, G62132, G62151, G62153, G63153, and Gajah) have stable starch content. G61151 has the highest rank of starch content estimation stability (21.22%). Spearman's correlation demonstrated the relationship of various methods of static or dynamic stability of starch content estimation (%).

Starch content estimation (%) has highly significant and positive correlation with Top, NP_i^2 , NP_i^3 , and NP_i^4 , but is negatively correlated with RS (Table 5). The stability values were only obtained by Huehn ($S1^{(1)}$, $S1^{(2)}$, $S1^{(3)}$, $S1^{(6)}$) method and did not have correlation with Thennarasu's (NP_i^1 , NP_i^2 , NP_i^3 , NP_i^4) method.

These results were similar to those reported by Rahadi et al. (2013), but are different from Sabaghnia et al. (2012) who reported that Thennarasu method had a high correlation with yield.

Principal component analysis (PCA) was described by Spearman's correlation. Zulhayana (2010) showed the high and positive value of nonparametric stability index identified the dynamic stability genotype. The result of principal component analysis (PCA) in Figure 2 divides the nonparametric stability methods into three groups, C1, C2, and C3.

Top, NP_i^2 , NP_i^3 , and NP_i^4 parameter and average starch content estimation (Y) are in the same group (C1), $S1^{(1)}$, $S1^{(2)}$, $S1^{(3)}$, $S1^{(6)}$, and NP_i^2 parameters are in group C2, whereas RS (Rank-Sum) is in group C3. Group C1 shows the concept of dynamic stability because Top, NP_i^2 , NP_i^3 , NP_i^4 method has the highest and positive correlation with the starch content estimation. Mut et al. (2008) showed that Top and RS methods were included as dynamic stability, however RS in this study has the negative correlation. The

Table 5. Spearman correlation of nonparametric stability parameter against the starch content estimation (%) of 16 genotypes in three environments

	SCE (%)	$S1^{(1)}$	$S1^{(2)}$	$S1^{(3)}$	$SI^{(6)}$	TOP	RS	NP_i^1	NP_i^2	NP_i^3
$S1^{(1)}$	-0.170 0.530									
$S1^{(2)}$	0.133 0.624	0.770**								
$S1^{(3)}$	0.142 0.601	0.754**	0.905**							
$SI^{(6)}$	0.242 0.366	0.783**	0.895**	0.957**						
TOP	0.703** 0.002	-0.122	-0.017	0.035	0.151					
RS	-0.515* 0.041	0.585*	0.486	0.365	0.317	-0.592*				
NP_i^1	0.058 0.831	0.718**	0.708**	0.620*	0.688**	0.013	0.713**			
NP_i^2	0.712** 0.002	0.213	0.235	0.134	0.261	0.541*	-0.027	0.450		
NP_i^3	0.793** 0.000	0.222	0.377	0.315	0.456	0.528*	0.020	0.604*	0.863**	
NP_i^4	0.790** 0.000	0.216	0.367	0.316	0.454	0.516*	0.019	0.597*	0.832**	0.995**
		0.423	0.162	0.232	0.077	0.041	0.944	0.015	0.000	0.000

Note: SCE= starch content estimation (%), * and **= significant at the levels of 5% and 1%; $S1^{(1)}$, $S1^{(2)}$, $S1^{(3)}$, $S1^{(6)}$ (Nassar and Huehn, 1987); Top (Fox et al., 1990); RS= Rank-sum by Kang (1988); NP_i^1 , NP_i^2 , NP_i^3 , NP_i^4 (Thennarasu, 1995).




Figure 2. Principal component analysis of starch content estimation (Y) and 10 nonparametric stability methods in cassava genotypes in three environments, C1, C2, and C3.

Table 6. Ranking of starch content estimation (Y) and nonparametric stability of 16 genotypes in 3 environments

Genotype	Dynamic stability					Frequency	Static stability					Frequency	
	Y	Top	NPi ²	NPi ³	NPi ⁴		S1 ¹	S1 ²	S1 ³	SI ⁶	RS		NPi ¹
G61122	13	12	10	9	9	0	15	12	10	10	16	12	0
G61142	2	2	11	7	8	4	1	5	6	6	1	1	5
G61143	14	12	12	3	4	0	5	3	2	3	10	6	3
G61151	12	12	2	2	2	5	7	7	7	7	5	4	1
G62143	10	12	5	12	12	0	13	16	15	15	16	15	0
G62132	16	7	3	4	3	4	9	6	5	5	12	13	1
G62151	4	2	15	14	14	0	11	8	8	8	3	7	1
G62153	15	12	1	1	1	5	6	1	1	1	6	2	6
G63111	11	12	4	5	5	4	14	13	14	12	6	8	1
G63121	7	7	6	10	10	2	4	11	9	11	6	9	1
G63124	5	7	8	8	7	2	3	2	3	2	2	3	6
G63142	1	2	16	16	16	0	12	14	13	14	7	16	0
G63143	8	7	7	6	6	2	10	10	11	9	6	5	1
G63153	9	7	9	11	11	2	16	15	16	16	10	14	0
“Gajah”	6	12	14	13	13	0	2	4	4	4	10	10	3
“Manggu”	3	7	13	15	16	0	8	9	12	13	4	11	1

Note: Y= starch content estimation (%); S1¹, S1², S1³, S1⁶ (Nassar and Huehn, 1987); Top, Mid, and Low by Fox et al. (1990); RS by Kang (1988); NPi¹, NPi², NPi³, and NPi⁴ (Thennarasu, 1995).

effect of high environmental diversity caused the changes in stability value.

Static stability concept can be used to select the best genotype with the highest production in different environmental conditions, the stability of genotype was also determined by ranking frequency (Table 6). Ranking frequency of two stability methods (dynamic and static) shows the information about genotype stability with the starch content estimation (%).

Static and dynamic stability concepts depend on range of region and testing sites, the more extensive the range of testing sites the more diversity in site conditions. Table 6 shows the top five ranks based on starch content estimation (%) are G61142, G62151, G63124, G63142, and "Manggu". The nine genotypes categorized as statically stable (yield stable in each environment) are G61151, G62153, G61142, G62132, G63111, G63121, G63124, G63143, and G63153. The static stability concept is highly dependent on the range of region and testing sites.

Twelve cassava genotypes are categorized as dynamically stable (adaptation followed the environmental index); they are G62153, G63124, G61142, G61143, Gajah, G61151, G62132, G62151, G63111, G63121, G63143, and "Manggu". The dynamic stability concept is more appropriate as the basis of selecting genotypes at the time of releasing genotype varieties because can explain the stability and adaptability of a genotype. G61142 was categorized as statically and dynamically stable, the most stable genotype and potential to released as new variety. This is related to the purpose of releasing varieties that still need improvement in starch content estimation quantity.

Conclusion

Variant analysis of the combined 16 cassava genotypes tested in three environments showed that the interaction between genotype and location did not have significant effects in each environment. Cassava genotypes, however, demonstrated different starch content estimation measured by gravimetric method. G63142 had the highest starch content estimation of 29.99%, whereas the top five genotypes based on the percentage of starch content estimation are G63142, G61142, local variety "Manggu", G62151, and G63124. G61142 was categorized as statically and dynamically stable, the most stable genotype, and has potentials to be released as a new variety. G63124 was categorized as statically stable, whereas G63142, G62151, and local "Manggu" had the highest diversity and high

percentage of starch content estimation.

Acknowledgement

The authors thanked The Ministry of Research, Technology and Higher Education, Republic of Indonesia, for the funding of this research through KLN-PI and Research-based Competence Scheme for the late Dr. Nurul Khumaida.

References

- Akcura, M., Taner, K.S., and Ayranci, R. (2006). Parametric stability analyses for grain yield of durum wheat. *Plant Soil and Environment* **6**, 254-261.
- Anasari, N.R., Kendarini, N., and Purnamaningsih, S.L. (2017). Genotype x environment interaction of four genotype of pakchoy (*Brassica rapa* L.) at three location. *Produksi Tanaman* **5**, 54-60.
- [BPS] Badan Pusat Statistik. (2017). "Ekspor-Impor Pati Ubi Kayu 2017". <http://www.bps.go.id>. [December 13, 2017].
- Baye, T.M., Abebe, T., and Wilke, R.A. (2011). Genotype environment interactions and their translational implications. *Personalized Medicine* **8**, 59-70.
- Chahal, G.S., and Gosal, S.S. (2002). "Principles and Procedures of Plant Breeding Biotechnological and Conventional Approaches". Narosa Publishing House. New Delhi.
- De Vita, P., Mastrangelo, A.M., Matteu, L., Mazzucotelli, E., Virzi, N., Palumbo, M., Lo Storto, M., Rizza, F., and Cattaveli, L. (2010). Genetic improvement effects on yield stability in durum wheat genotypes grown in Italy. *Field Crop Research* **199**, 68-77.
- Fahreza, A. (2014). "Karakterisasi Morfologi dan Pertumbuhan Beberapa Genotipe Ubi Kayu (*Manihot esculenta* Crantz) Hasil Induksi Mutasi menggunakan Iradiasi Sinar Gamma". [Thesis]. Bogor Agricultural University.
- [FAO] Food and Agriculture Organization. (2017). "Food Outlook, Biannual Report on Global Food Markets". <http://www.fao.org/giews/english/fo/index.html> [December 13, 2017].
- Fox, P.N., Skovmand, B.K., Braun, H.J., and Comier,

- R. (1990). Yield and adaptation of hexaploid spring triticale. *International Journal of Plant Breeding* **47**, 57-64.
- Haloho, J.D. (2014). "Pengolahan Ubi Kayu dalam Upaya Percepatan Diversifikasi Pangan di Kalimantan Barat". pp 775-786. Balitkabi. Malang. Indonesia.
- Harahap, Z. (1982). "Pedoman Pemuliaan Padi". Lembaga Biologi Nasional. Indonesia.
- Huehn, M. (1990). Nonparametric measures of phenotypic stability. *International Journal of Plant Breeding* **7**, 189-194.
- Kang, M.S. (1988). A rank-sum method for selecting high yielding, stable corn genotypes. *Cereal Research Communication* **16**, 113-115.
- Kawano, K., Fukuda, W.M.G., and Cempukdee, U. (1987). Genetic and environmental effect on dry matter content of cassava root. *Crop Science* **27**, 69-74.
- Lebot, V. (2009). "Tropical Root and Tuber Crops : Cassava, Sweet Potato, Yams, and Aroids". CABI. United Kingdom.
- Maharani, S. (2015). "Iradiasi Sinar Gamma pada Lima Genotipe Ubi Kayu (*Manihot esculenta* Crantz.) dan Pengujian Awal Stabilitas Mutan". [Thesis]. Bogor Agricultural University.
- Mohammadi, R., Roostaei, M., Ansari, Y., Aghaee, M., and Amri, A. (2010). Relationships of phenotypic stability measures for genotypes of three cereal crops. *Canadian Journal of Plant Science* **90**, 819-930.
- Murtiningrum, Bosawer, E.F., Istalaksana, P., and Jading, A. (2012). Karakterisasi umbi dan pati lima kultivar ubi kayu (*Manihot esculenta*). *Scientific Journal of Agricultural Science* **3**, 81-90.
- Mut, Z., Aydin, N., Bayramoglu, H.O., and Ozcan, H. (2009). Interpreting genotype x environment interaction in bread wheat (*Triticum aestivum* L.) genotypes using nonparametric measures. *Turkish Journal of Agriculture and Forestry* **33**, 127-137.
- Nassar, R.F., and Huehn, M. (1987). Studies on estimation on phenotypic stability: tests of significance for nonparametric measures of phenotypic stability. *Biometrics* **43**, 45-53.
- [PERMENTAN] Peraturan Kementerian Pertanian. (2011). "Varietas unggul". <http://perundangan.pertanian.go.id/admin/file/Permentan-70-11> [December 13, 2017].
- Rahadi, V.P., Syukur, M., Sujiprihati, S., and Yuniarti, R. (2013). Nonparametric stability analysis of yield for nine chili pepper (*Capsicum annuum* L.) genotypes in eight environments. *AGRIVITA : Journal of Agricultural Science* **35**, 193-200.
- Sabaghnia, N., Karimizadeh, R., and Mohammadi, M. (2012). The use of corrected and uncorrected nonparametric stability measurements in durum wheat multi-environments trial. *Spanish Journal of Agricultural Research* **10**, 722-730.
- Satoto, S., Utomo, T.W., Widyastuti, Y., and Rumanti, I.A. (2009). "Submission of release for new hybrid rice variety H45, H47, H64, H68, and H78. [Paper of Variety Release on Trial]". pp 20. Balai Besar Penelitian Padi. Subang. Indonesia.
- Subekti, I. (2013). "Karakterisasi Morfologi dan Pertumbuhan Ubi Kayu 'Gajah' Asal Kalimantan Timur Hasil Iradiasi Sinar Gamma". [Thesis]. Bogor Agricultural University.
- Subekti, I., Khumaida, N., Ardie, S.W., and Syukur, M. (2018). Evaluasi hasil dan kandungan pati mutan ubi kayu hasil iradiasi sinar gamma generasi M₁V₄. *Indonesian Journal of Agronomy* **46**, 64-70.
- Syukur, M., Sujiprihati, S., and Yuniarti, R. (2012). "Teknik Pemuliaan Tanaman". Penebar Swadaya. Indonesia.
- Thennarasu, K. (1995). "On certain nonparametric procedures for studying genotype-environment interactions and yield stability". P.J. School, New Delhi.
- Zulhayana, S. (2010). "Klasifikasi Genotipe dengan Pendekatan Indeks Stabilitas Nonparametrik" [Thesis]. Bogor Agricultural University.